


Section 3.2 – Inverse Functions and Logs

Inverse Relations:

- To find an _____ you are looking to 'undo' the process that was done with the original relation.
- For example, the inverse of the function $f(x) = \frac{1}{x-2}$ is the function $g(x) = \frac{1}{x} + 2$.



Notice that $f(3) = 1$, and $g(1) = 3$.


- Also notice in the previous example, we have the ordered pair (3,1) for $f(x)$ and the ordered pair (1,3) for $g(x)$. In fact, being able to interchange the first and second coordinates of each ordered pair in a relation is another way to recognize an _____.
- To find an inverse relation for $y = \text{relation of } x$, interchange the x 's and y 's in the equation and (if possible) solve for y .
- *Example. Find the equation of the inverse of $y = x^2 - 6x + 9$.*

- **Q:** Is the original problem a function? Does it pass the vertical line test? Is the inverse a function?
A: _____

Inverse Functions and One-to-One:

- The concepts above hold for functions as well. You can find an inverse function by switching the order of all ordered pairs, or you can switch the x 's and y 's in the equation and solve.
- The notation for inverse functions is a little bit different, because for a function $f(x)$, whose inverse is also a function, the notation for the inverse is $f^{-1}(x)$.
- Note, this does NOT mean $\frac{1}{f(x)}$, but f inverse!

- Notice in the problem given previously, $f(x) = x^2 - 6x + 9$ is a function. But the inverse, $\pm\sqrt{x} + 3$




is NOT a function.

- Q: How can you tell the inverse is not a function?**

A: _____.

- Recall that for $y(x)$ to be a function, each x has only one y .
- $y(x)$ is _____ if it is
 - a function, and
 - each y maps back to only one x
- For one-to-one, each x has only one y , and each y has only one x . It has to pass not only the vertical line test, but also a _____.
- Q: Which of the following are functions? Which are one-to-one?**



A: _____.

- The following functions are always one-to-one:
Linear, square roots
- The following functions are not ever one-to-one:
Quadratic, absolute value

- *Example.* Graph $f(x) = \frac{5x-3}{2x+1}$, determine if it is 1-1, if so find the inverse




- **Q:** What is the relationship between the domain and ranges for functions and their inverses?

A: _____.

- Since an inverse function ‘undoes’ the original function, if we compose the two [i.e. take $f \circ f^{-1}(x)$ or $f^{-1} \circ f(x)$] we will get x .

- *Example.* Show $f(x) = \frac{x+5}{4}$, $f^{-1}(x) = 4x-5$ are inverses



Log Functions:

- Recall: $b^y = x$ if and only if $y = \log_b x$
 y is the exponent, b is the base, and x is the argument
- **Special Log Bases:**
Log base e is _____ (written \ln)
Log base 10 is _____ (written \log)

- **Log Rules:**

$$\log_b b = 1$$

$$\log_b 1 = 0$$

$$\log_b b^p = p$$

$$\log_b (MN) = \log_b M + \log_b N$$

$$\log_b M^p = p \log_b M$$

$$\log_b (M / N) = \log_b M - \log_b N$$

$$b^{\log_b p} = p$$

- *Example. Express $\log_a \sqrt{\frac{a^6 b^8}{a^2 b^5}}$ in terms of sums and differences*

- *Example, simplify $2\log_5 x - \log_5 y - 3\log_5 z$*

Solving Exponential Equations:

- Equations with variables in the exponent are called _____.
- If the base is the same on both sides of the equation, you can equate the exponents.

- *Example. Solve $3^{x^2+4x} = \frac{1}{27}$*

- More work comes when the base is not the same. You will then have to solve the equation using logs.

- *Example. Solve $2^x = 40$*

- *Example. Solve $e^x - 6e^{-x} = 1$*

Solving Log Equations:

- Often it is useful to change to exponential in form.
- *Example. Solve $\log_2 x = -3$*

- *Example. Solve $\log_5(8 - 7x) = 3$*

- *Example. Solve $\log x - \log(x + 3) = -1$*